arXiv:1112.6347v1 [math.GR] 29 Dec 2011

Alternating subgroups of Coxeter groups and their spinor extensions

O. V. Ogievetsky$^{\circ\diamond 1}$ **and L. Poulain d'Andecy**$^{\circ}$

∘ Center of Theoretical Physics, Luminy
13288 Marseille, France

⋄ J.-V. Poncelet French-Russian Laboratory, UMI 2615 du CNRS, Independent University of Moscow, 11 B. Vlasievski per., 119002 Moscow, Russia

Abstract

Let G be a discrete Coxeter group, G^+ its alternating subgroup and $\tilde{G}^+$ the spinor cover of G^+. A presentation of the groups G^+ and $\tilde{G}^+$ is proved for an arbitrary Coxeter system (G, S); the generators are related to edges of the Coxeter graph. Results of the Coxeter–Todd algorithm - with this presentation - for the chains of alternating groups of types A, B and D are given.

1. Introduction

Let (G, S) be a Coxeter system and G^+ the alternating subgroup of G. A presentation of G^+ by generators and relations is given, as an exercise, in [1]. One (arbitrary) vertex in the Coxeter graph is distinguished in this presentation.

For the Coxeter groups of type A (that is, symmetric groups) the standard Artin presentation equips the chain of symmetric groups with a structure of a local and stationary chain of groups. The representation theory of the chain of symmetric groups was revisited in [8]. It was emphasized in [8] that the method developed there for the representation theory of the symmetric groups could be hopefully relevant for a wider class of local and stationary chains of groups or algebras (see [11] for the definition of locality and stationarity). For type A, the Bourbaki presentation of the chain of the alternating groups is not local.

A discrete Coxeter group G is a subgroup of an orthogonal group O; the spin cover of O restricts to a central extension $\tilde{G}$ of G. A presentation of $\tilde{G}$ can be found in [7]. The subgroup G^+ of a discrete Coxeter group G inherits as well a central (spinor) extension $\tilde{G}^+$. In general, there are central extensions of alternating groups different from the spinor ones (see [6] for Schur multipliers of alternating subgroups of finite Coxeter groups and for a presentation à la Bourbaki).

We suggest another presentation of the groups G^+ and $\tilde{G}^+$ for an arbitrary Coxeter system (G, S). In the Bourbaki presentation the generators are indexed by vertices (except the distinguished one) of the Coxeter graph. In our presentation the generators are indexed by edges of the Coxeter graph.

---

[1] On leave of absence from P. N. Lebedev Physical Institute, Leninsky Pr. 53, 117924 Moscow, Russia

For type A our presentations of the chains of the groups A_n^+ and $\tilde{A}_n^+$ are local and stationary. For the chain of the groups A_n^+ a similar (but not exactly the same) presentation was given in [12] and it was asked there how to generalize this presentation to all alternating groups. For types B and D, our presentations of the chains of groups B_n^+, $\tilde{B}_n^+$, D_n^+ and $\tilde{D}_n^+$ are local and eventually stationary (that is, stationary for $n \geqslant n_0$ for a certain n_0).

The group A_n^+ has a presentation referred to as *Carmichael presentation* in [3]. We interpret this presentation geometrically: here one distinguishes an oriented edge in the Coxeter graph. With this interpretation there are analogues of the Carmichael generators for other Coxeter systems. In general, these elements generate a subgroup of G^+. However for type B (with an appropriate choice of a distinguished oriented edge) and type D the whole group G^+ is obtained. We give defining relations for these sets of generators of B_n^+ and D_n^+.

The paper is organized as follows. In Section 2 we obtain the presentation of the alternating subgroup G^+ for a Coxeter system (G, S). We first deal with irreducible Coxeter systems. Then we generalize the presentation to arbitrary Coxeter systems by introducing a *connected extension* of the Coxeter graph. Section 3 is devoted to the presentation of the group $\tilde{G}^+$ for an arbitrary Coxeter system (G, S). One can give presentations - with generators related to edges of the Coxeter graph - for all central extensions of the alternating subgroups of finite Coxeter groups. We illustrate it on two examples: A_5^+ and A_6^+. In Section 4 we give results of the Coxeter–Todd algorithm for three presentations - the Bourbaki one, our presentation which refers to edges and the presentation à la Carmichael - of the alternating groups of types A, B and D.

In the text whenever a product $\pi\sigma$ of two permutations appears we assume that σ is applied first and then π; for example, $(1,2)(2,3) = (1,2,3)$.

2. Alternating subgroups of Coxeter groups

1. Let (G, S) be a Coxeter system: S is the set of generators, $S = \{s_0, \ldots, s_{n-1}\}$; the defining relations are encoded by a symmetric matrix $\mathfrak{m} = (m_{ij})_{i,j=0,1,\ldots,n-1}$ with $m_{ii} = 1$ and $2 \leq m_{ij} \in \mathbb{N} \cup \infty$:

$$G := \langle\, s_0, \ldots, s_{n-1} \mid (s_i s_j)^{m_{ij}} = 1 \text{ for } i, j = 0, 1, \ldots, n-1,\ i \leqslant j \,\rangle\,. \tag{1}$$

This presentation of G can be expressed with the help of a *Coxeter* graph: its vertices are in one-to-one correspondence with the generators $s_0, \ldots, s_{n-1}$ and are indexed by the subscripts $0, 1, \ldots, n-1$; the vertex i is connected to the vertex j if and only if $m_{ij} \geq 3$; the edge between i and j is labeled by the number m_{ij} if $m_{ij} \geq 3$ (for brevity, in the sequel we draw a simple line for an edge labeled by 3 and a double line for an edge labeled by 4). We denote the Coxeter graph corresponding to the Coxeter system (G, S) by $\mathcal{G}$.

The sign character is the unique homomorphism $\epsilon : G \to \{-1, 1\}$ such that $\epsilon(s_i) = -1$ for $i = 0, \ldots, n-1$. Its kernel $G^+ := \ker(\epsilon)$ is called the alternating subgroup of G. Recall the Bourbaki [1] presentation of G^+, see [2] for a proof. The alternating group G^+ is isomorphic to the group generated by $R_1, \ldots, R_{n-1}$ with the defining relations:

$$\begin{cases} R_i^{m_{0i}} = 1 & \text{for } i = 1, \ldots, n-1, \\ (R_i^{-1} R_j)^{m_{ij}} = 1 & \text{for } i, j = 1, \ldots, n-1 \text{ such that } i < j. \end{cases} \tag{2}$$

The isomorphism with G^+ is given by $R_i \mapsto s_0 s_i$ for $i = 1, \dots, n-1$. The Bourbaki presentation of the alternating group G^+ depends on the choice of a generator carrying the subscript 0.

2. In this paragraph, we assume that the Coxeter system (G, S) is irreducible (that is, the Coxeter graph $\mathcal{G}$ is connected). We give a presentation of G^+ in terms of generators corresponding to the edges of the Coxeter graph; no vertex in the Coxeter graph is distinguished. In the next paragraph this presentation will be generalized to arbitrary Coxeter systems.

The presentation uses an orientation - chosen arbitrarily - of edges of the Coxeter graph. For concreteness, if there is an edge between i and j with $i < j$, we orient it from i to j. We associate a generator r_{ij} to every oriented edge, that is, to every pair (i, j), $i, j = 0, \dots, n-1$, such that $i < j$ and $m_{ij} \neq 2$. For a generator r_{ij} we denote by r_{ji} the inverse, $r_{ji} := r_{ij}^{-1}$.

Definition 2.1 *Two generators r_{ij} and r_{lm} are said to be not connected if $\{i, j\} \cap \{l, m\} = \varnothing$ and there is no edge connecting any of the vertices $\{i, j\}$ with any of the vertices $\{l, m\}$.*

Proposition 2.2 *Let (G, S) be an irreducible Coxeter system with a Coxeter matrix $\mathfrak{m}$. The alternating subgroup G^+ of G is isomorphic to the group with generators r_{ij} and the defining relations*

$$\begin{cases} (r_{ij})^{m_{ij}} = 1 & \textit{for all generators } r_{ij}, & (3) \\ r_{ii_1} r_{i_1 i_2} \dots r_{i_a i} = 1 & \textit{for cycles with edges } (ii_1), (i_1 i_2) \dots, (i_a i), & (4) \\ (r_{ij} r_{jk})^2 = 1 & \textit{for } r_{ij}, r_{jk} \textit{ such that } i < k \textit{ and } m_{ik} = 2, & (5) \\ (r_{ij} r_{jk} r_{kl})^2 = 1 & \textit{for } r_{ij}, r_{jk}, r_{kl} \textit{ such that } i < l \textit{ and } m_{il} = 2, & (6) \\ r_{ij} r_{lm} = r_{lm} r_{ij} & \textit{for } r_{ij} \textit{ and } r_{lm} \textit{ which are not connected.} & (7) \end{cases}$$

Let $\mathfrak{c}_1, \dots, \mathfrak{c}_{\mathfrak{l}}$ be a set of generators of the fundamental group of $\mathcal{G}$. In the set of the defining relations it is sufficient to impose the relation (4) for the cycles $\mathfrak{c}_{\mathfrak{a}}$, $\mathfrak{a} = 1, \dots, \mathfrak{l}$. If G is finite, its Coxeter graph has no cycles so the relation (4) is not needed.

Proof of the Proposition. Let W^+ be the group generated by r_{ij} with the defining relations (3)–(7). Define the map ϕ from the set of generators r_{ij} of W^+ to G^+ by

$$\phi \ : \ r_{ij} \mapsto \begin{cases} R_i^{-1} R_j & \text{if } i \neq 0, \\ R_j & \text{if } i = 0. \end{cases} \tag{8}$$

It is straightforward to see that the relations (3)–(6) are verified by the images of the generators r_{ij} in G^+. As for the last one, the generators r_{ij} and r_{lm} are not connected, so $m_{il} = m_{im} = m_{jl} = m_{jm} = 2$. We have two possibilities: either $i \neq 0$ and $l \neq 0$ or one of the numbers, i or l, is 0.

- If $i \neq 0$ and $l \neq 0$ then

$$R_i^{-1} R_j R_l^{-1} R_m = R_i^{-1} R_l R_j^{-1} R_m = R_l^{-1} R_i R_j^{-1} R_m = R_l^{-1} R_i R_m^{-1} R_j = R_l^{-1} R_m R_i^{-1} R_l, \tag{9}$$

  where we have repeatedly used the second line in (2). Thus $\phi(r_{ij}) \phi(r_{lm}) = \phi(r_{lm}) \phi(r_{ij})$.

- If, say, $l = 0$ then $i \neq 0$ and we have $R_i^2 = R_j^2 = 1$ because $m_{0i} = m_{0j} = 2$. We calculate

  $$R_i^{-1} R_j R_m = R_i^{-1} R_m^{-1} R_j = R_m R_i^{-1} R_j; \tag{10}$$

  we used $(R_j^{-1} R_m)^2 = 1$ and $R_j^2 = 1$ in the first equality; in the second equality we used $(R_i^{-1} R_m)^2 = 1$ and $R_i^2 = 1$. Thus $\phi(r_{ij})\phi(r_{lm}) = \phi(r_{lm})\phi(r_{ij})$ again.

We conclude that ϕ extends to a group homomorphism from W^+ to G^+.

Define now the map ψ from the set of generators of G^+ to W^+ by

$$\psi \ : \ R_i \mapsto \grave{R}_i := r_{0i_1} r_{i_1 i_2} \ldots r_{i_k i} \ \text{ for all } i = 1, \ldots, n-1, \tag{11}$$

where $(0, i_1, i_2, \ldots, i_k, i)$ is an arbitrary path from the vertex 0 to the vertex i in the Coxeter graph (such path exists because the Coxeter graph is connected). Due to the relation (4), the element $\grave{R}_i$ does not depend on the chosen path, thus the map ψ is well-defined. We shall assume that $(0, i_1, i_2, \ldots, i_k, i)$ is a shortest (that is, with minimal number of edges) path from 0 to i.

If $j \neq i$ then

$$\grave{R}_i^{-1} \grave{R}_j = r_{ii_k} \ldots r_{i_1 0} r_{0 j_1} \ldots r_{j_l j} \ , \text{ or, again by (4), } \grave{R}_i^{-1} \grave{R}_j = r_{ii_a} \ldots r_{i_b j},$$

where $(i, i_a, \ldots, i_b, j)$ is a shortest path from i to j.

We shall verify that $\grave{R}_i^{m_{0i}} = 1$ for $i = 1, \ldots, n-1$, and $(\grave{R}_i^{-1} \grave{R}_j)^{m_{ij}} = 1$ for $i, j = 1, \ldots, n-1$ with $i < j$. It suffices to prove that $(r_{ia_1} r_{a_1 a_2} \ldots r_{a_k j})^{m_{ij}} = 1$ for a shortest path $(i, a_1, a_2, \ldots, a_k, j)$. If $(k_1, k_2, \ldots, k_l)$ is a shortest path then there is no edge in the Coxeter graph between the vertices k_μ and k_ν with $|\mu - \nu| > 1$. If $m_{ij} \neq 2$ the shortest path is (i, j) and $(r_{ij})^{m_{ij}} = 1$. If $m_{ij} = 2$ and $k = 1$ (respectively, $k = 2$) then $(r_{ia_1} \ldots r_{a_k j})^2 = 1$ by (5) (respectively, (6)). If $m_{ij} = 2$ and $k \geq 3$ we have:

$$\begin{aligned}(r_{ia_1} r_{a_1 a_2} \ldots r_{a_k j})^2 \quad &= r_{ia_1} r_{a_1 a_2} r_{a_2 a_3} \ldots r_{a_k j} r_{ia_1} \ldots r_{a_k j} = r_{a_2 a_3}^{-1} r_{a_1 a_2}^{-1} r_{ia_1}^{-1} \ldots r_{a_k j} r_{ia_1} \ldots r_{a_k j} \\ &= r_{a_1 a_2} r_{a_2 a_3} \ldots r_{a_k j} r_{a_1 a_2} \ldots r_{a_k j} = (r_{a_1 a_2} \ldots r_{a_k j})^2.\end{aligned}$$

In the second equality we used that $(r_{ia_1} r_{a_1 a_2} r_{a_2 a_3})^2 = 1$ since $m_{ia_3} = 2$; in the third equality, we used that r_{ia_1} commutes with $r_{a_3 a_4} \ldots r_{a_k j}$ and, since $m_{a_1 a_3} = 2$, that $(r_{a_1 a_2} r_{a_2 a_3})^2 = 1$. These properties are implied by the the minimality of the path $(i, a_1, a_2, \ldots, a_k, j)$. The assertion follows by induction on the length of the path.

We conclude that the map ψ extends to a group homomorphism from G^+ to W^+. It is straighforward to see that the morphisms ϕ and ψ are mutually inverse. $\square$

Remark. Let $(i, a_1, a_2, \ldots, a_k, j)$ be any path in the Coxeter graph. As the proof shows, the relations (3)–(7) imply $(r_{ia_1} r_{a_1 a_2} \ldots r_{a_k j})^{m_{ij}} = 1$. Therefore, the defining relations (3)–(7) are equivalent to the following relations:

$$\begin{cases} (r_{ia_1} r_{a_1 a_2} \ldots r_{a_k j})^{m_{ij}} = 1 & \text{for any path } (i, a_1, a_2, \ldots, a_k, j) \text{ in the Coxeter graph,} \\ r_{ij} r_{lm} = r_{lm} r_{ij} & \text{for } r_{ij} \text{ and } r_{lm} \text{ which are not connected.} \end{cases}$$

3. Now assume that the Coxeter graph $\mathcal{G}$ is not connected. We add some 'virtual" edges labeled by 2 to make the graph connected. More precisely, let $\mathcal{G} = \mathcal{G}_1 \sqcup \mathcal{G}_2 \sqcup \cdots \sqcup \mathcal{G}_m$ be a decomposition of $\mathcal{G}$ into the disjoint union of its connected components. For $a = 1, \ldots, m$ choose an arbitrary vertex i_a of $\mathcal{G}_a$; then, for all $l = 1, \ldots, m-1$, we add a virtual edge between i_l and i_{l+1} and label it by 2. Virtual edges will be drawn with dashed lines. The obtained graph we call a *connected extension* of $\mathcal{G}$.

A connected extension is not unique. Fix a connected extension $\mathcal{G}^c$ of $\mathcal{G}$ and associate a generator r_{ij}, $i < j$, to every oriented edge (virtual or not) of $\mathcal{G}^c$. The analogue of the Definition 2.1 includes now virtual edges.

Definition 2.3 *Two generators r_{ij} and r_{kl} are said to be not connected if $\{i, j\} \cap \{l, m\} = \varnothing$ and there is no edge or virtual edge connecting any of the vertices $\{i, j\}$ with any of the vertices $\{l, m\}$.*

With this choice of the set of generators, the formulation of the Proposition 2.2 stays the same for an arbitrary Coxeter system.

Proposition 2.4 *Let (G, S) be a Coxeter system with a Coxeter matrix $\mathfrak{m}$. The alternating subgroup G^+ of G is isomorphic to the group generated by r_{ij} with the defining relations (3)–(7).*

Proof. The proof goes exactly as the proof of the Proposition 2.2. The isomorphisms are still given by

$$\phi \ : \ r_{ij} \mapsto \begin{cases} R_i^{-1} R_j & \text{if } i \neq 0, \\ R_j & \text{if } i = 0, \end{cases} \tag{12}$$

and

$$\psi \ : \ R_i \mapsto r_{0i_1} r_{i_1 i_2} \ldots r_{i_k i} \ \text{ for } i = 1, \ldots, n-1, \tag{13}$$

where $(0, i_1, i_2, \ldots, i_k, i)$ is an arbitrary path from the vertex 0 to the vertex i in $\mathcal{G}^c$. □

Remark. Denote by $G(\mathcal{G})$ the Coxeter group related to the Coxeter graph $\mathcal{G}$. The defining relations (3)–(7) imply that generators r_{ij} and r_{kl} commute if the edges (i, j) and (k, l) belong to different connected components of $\mathcal{G}$. Indeed, either r_{ij} and r_{kl} are not connected and they commute, or there is a virtual edge between, say, vertices j and k; in the latter situation, relations (5)–(6) gives

$$r_{kl} r_{ij} = r_{jk}^{-1} r_{ij}^{-1} r_{kl}^{-1} r_{jk}^{-1} = r_{ij} r_{jk}^2 r_{kl},$$

and the assertion follows from $r_{jk}^2 = 1$.

Let, as above, $\mathcal{G} = \mathcal{G}_1 \sqcup \mathcal{G}_2 \sqcup \cdots \sqcup \mathcal{G}_m$; the groups $G(\mathcal{G}_a)$, $a = 1, \ldots, m$, are naturally subgroups of $G(\mathcal{G})$. The virtual edges form a path in $\mathcal{G}^c$ and one can verify that the generators r_{st} corresponding to the virtual edges of $\mathcal{G}^c$ generate a subgroup $\mathcal{Y}$ isomorphic to C_2^{m-1}, where C_2 is the cyclic group with 2 elements. One can also verify that each $G(\mathcal{G}_a)$, $a = 1, \ldots, m$, is stable under conjugations by elements of $\mathcal{Y}$. Thus, each $G(\mathcal{G}_a)$, $a = 1, \ldots, m$, is normal in $G(\mathcal{G})$ and $G(\mathcal{G})$ is the semi-direct product $\mathcal{Y} \ltimes (G(\mathcal{G}_1) \times G(\mathcal{G}_2) \times \cdots \times G(\mathcal{G}_m))$.

Example. Consider the Coxeter group generated by s_1, s_2, s_3, s_4, s_5 with the Coxeter matrix $\mathfrak{m} = \begin{pmatrix} 1 & 4 & 2 & 2 & 2 \\ 4 & 1 & 2 & 2 & 2 \\ 2 & 2 & 1 & 3 & 3 \\ 2 & 2 & 3 & 1 & 3 \\ 2 & 2 & 3 & 3 & 1 \end{pmatrix}$. The Coxeter graph has two components which we connect by adding a virtual edge between vertices 2 and 3. The figure shows oriented edges and generators corresponding to each oriented edge and virtual edge. With the Definition 2.3 there is only one pair of generators which are not connected: r_{12} and r_{45}.



Fig. 1. *Example of a connected extension of the Coxeter graph*

The defining relations for the generators r_{12}, r_{23}, r_{34}, r_{35} and r_{45} are:

$$\begin{cases} (r_{12})^4 = 1,\ (r_{34})^3 = 1,\ (r_{35})^3 = 1,\ (r_{45})^3 = 1,\ (r_{23})^2 = 1, \\ r_{34}r_{45}r_{53} = 1, \\ (r_{12}r_{23})^2 = 1,\ (r_{23}r_{34})^2 = 1,\ (r_{23}r_{35})^2 = 1, \\ (r_{12}r_{23}r_{34})^2 = 1,\ (r_{12}r_{23}r_{35})^2 = 1,\ (r_{23}r_{34}r_{45})^2 = 1,\ (r_{23}r_{35}r_{54})^2 = 1, \\ r_{12}r_{45} = r_{45}r_{12}. \end{cases}$$

4. We write explicitly the presentation of the Proposition 2.2 for the alternating groups of type A. The Coxeter group A_n is isomorphic to the group of permutations of $n+1$ elements (the symmetric group) and the alternating group A_n^+ is isomorphic to the subgroup of even permutations. We use the notation for the generators explained by the following figure (recall that a vertex i corresponds to a generator s_i of the Coxeter group and that a generator of the alternating group is associated to each oriented edge once an orientation is chosen; we have set $r_i := r_{i-1,i}$ for all $i = 1, \dots, n-1$):



Fig. 2. *Coxeter graph of type A*

According to the Proposition 2.2 the group A_n^+ is generated by r_i, $i = 1, \dots, n-1$, with the defining relations:

$$\begin{cases} {r_i}^3 = 1 & \text{for } i = 1, \dots, n-1, \\ (r_i r_{i+1})^2 = 1 & \text{for } i = 1, \dots, n-2, \\ (r_i r_{i+1} r_{i+2})^2 = 1 & \text{for } i = 1, \dots, n-3, \\ r_i r_j = r_j r_i & \text{for } i, j = 1, \dots, n-1 \text{ such that } |i-j| > 2. \end{cases} \tag{14}$$

The isomorphism with the group of even permutations of $n+1$ elements is established by $r_i \mapsto (i, i+1, i+2)$ where $(i, i+1, i+2)$ is the cyclic permutation of i, $i+1$ and $i+2$. In Section 4 we give a different proof, based on the Coxeter-Todd algorithm, for this presentation of A_n^+.

The presentation (14) is a local and stationary, in the sense of [11], presentation for the chain of groups A_n^+. In [12] a presentation of the group A_n^+ is proved. The generators are $\rho_1, \dots, \rho_{n-1}$ and the defining relations are the same as the defining relations (14) for the r_i, except that the third relation in (14) is replaced by $\rho_i\rho_{i+1}^2\rho_{i+2} = \rho_{i+2}\rho_i$ for $i = 1,\ldots,n-3$. The equivalence of the two presentations is given by $\rho_i \mapsto r_i$ for $i = 1,\ldots,n-1$. Indeed we have

$$(\rho_i\rho_{i+1}\rho_{i+2})^2 = \rho_i\rho_{i+1}\cdot\rho_{i+2}\rho_i\cdot\rho_{i+1}\rho_{i+2} = \rho_i\rho_{i+1}\cdot\rho_i\rho_{i+1}^2\rho_{i+2}\cdot\rho_{i+1}\rho_{i+2} = (\rho_i\rho_{i+1})^2(\rho_{i+1}\rho_{i+2})^2 = 1.$$

Conversely,

$$r_{i+2}r_i = r_{i+1}^{-1}r_i^{-1}\cdot r_{i+2}^{-1}r_{i+1}^{-1} = r_ir_{i+1}\cdot r_{i+1}r_{i+2} = r_ir_{i+1}^2r_{i+2}.$$

The Coxeter-Todd algorithm given in Section 4 provides a normal form, different from the normal form in [12], for the elements of the group A_n^+ .

Remark. The two first relations of (14) imply that:

$$\begin{cases} r_ir_{i+1}^2r_i = r_ir_{i+1}\cdot r_{i+1}r_i = r_{i+1}^2r_i^2\cdot r_i^2r_{i+1}^2 = r_{i+1}^2r_ir_{i+1}^2, \\ r_i^2r_{i+1}r_i^2 = r_i^2r_{i+1}^2\cdot r_{i+1}^2r_i^2 = r_{i+1}r_i\cdot r_ir_{i+1} = r_{i+1}r_i^2r_{i+1}, \end{cases} \qquad i = 1,\ldots,n-2\,.$$

Thus the generators $r_i' := r_i^{-1}$ for i odd and $r_i' := r_i$ for i even verify the Artin relation

$$r_i'r_{i+1}'r_i' = r_{i+1}'r_i'r_{i+1}' \quad \text{for } i = 1,\ldots,n-2.$$

3. Central extensions of the alternating subgroups of Coxeter groups

1. Let (G,S) be an arbitrary Coxeter system with $S = \{s_0,\ldots,s_{n-1}\}$ and the Coxeter matrix $\mathfrak{m}$. Let $\tilde{G}$ be the group with the generators $\tilde{s}_0,\ldots,\tilde{s}_{n-1}$, α and the defining relations

$$\begin{cases} (\tilde{s}_i\tilde{s}_j)^{m_{ij}} = 1 & \text{if } m_{ij} \text{ is odd,} \\ (\tilde{s}_i\tilde{s}_j)^{m_{ij}} = \alpha & \text{if } m_{ij} \text{ is even,} \\ \alpha \text{ is central and } \alpha^2 = 1. \end{cases} \tag{15}$$

Let also $\tilde{G}'$ be the group with the generators $\tilde{s}_0',\ldots,\tilde{s}_{n-1}'$, α' and the defining relations

$$\begin{cases} (\tilde{s}_i'\tilde{s}_j')^{m_{ij}} = \alpha', \\ \alpha' \text{ is central and } \alpha'^2 = 1. \end{cases} \tag{16}$$

When G is a discrete reflection group, $\tilde{G}$ and $\tilde{G}'$ are its spinor central extensions and they are non-trivial if there are some i,j, $i \neq j$, such that m_{ij} is even [7]. In general, the groups $\tilde{G}$ and $\tilde{G}'$ are not isomorphic. Note that discrete Coxeter groups admit non-trivial central extensions non-isomorphic to (15) or (16); see [5, 13] for the Schur multipliers of the discrete Coxeter groups.

Denote by $\pi\colon \tilde{G} \to G$ and $\pi'\colon \tilde{G}' \to G$ the natural projections. Let $\tilde{G}^+ := \pi^{-1}(G^+)$ and $\tilde{G}'^+ := \pi'^{-1}(G^+)$.

Our aim is to give presentations for the groups $\tilde{G}^+$ and $\tilde{G}'^+$ in the spirit of the preceding Section. As a by-product, we shall see that the groups $\tilde{G}^+$ and $\tilde{G}'^+$ are isomorphic.

We first give a presentation for $\tilde{G}^+$.

Proposition 3.1 *Let* (G, S) *be a Coxeter system with the Coxeter matrix* $\mathfrak{m}$*. The group* $\tilde{G}^+$ *is generated by* $\tilde{R}_1, \ldots, \tilde{R}_{n-1}$ *and* z *with the defining relations*

$$\begin{cases} (\tilde{R}_i)^{m_{0i}} = z^{m_{0i}-1} & \text{for } i = 1, \ldots, n-1, \qquad (17)\\ (\tilde{R}_i^{-1}\tilde{R}_j)^{m_{ij}} = z^{m_{ij}-1} & \text{for } i, j = 1, \ldots, n-1 \text{ such that } i < j, \qquad (18)\\ z \text{ is central and } z^2 = 1. & \qquad (19) \end{cases}$$

Proof. This is an adaptation of the proof, suggested in [1], of the presentation (2) for the group G^+; we give a sketch. Let $\tilde{W}^+$ be the group generated by the elements $\tilde{R}_1, \ldots, \tilde{R}_{n-1}$ and z with the defining relations (17)–(19). Define the map ϕ from the set of generators $\{\tilde{R}_1, \ldots, \tilde{R}_{n-1}, z\}$ to $\tilde{G}^+$ by:

$$\phi(\tilde{R}_i) = \tilde{s}_0\tilde{s}_i \ , \ i = 1, \ldots, n-1, \text{ and } \phi(z) = \alpha. \qquad (20)$$

This map extends to a surjective homomorphism from $\tilde{W}^+$ to $\tilde{G}^+$, which we denote by the same symbol ϕ. We will show that ϕ is an isomorphism.

One checks that the map ω given by $\omega(\tilde{R}_i) = \tilde{R}_i^{-1}$, $i = 1, \ldots, n-1$, and $\omega(z) = z$ extends to an involution of $\tilde{W}^+$, defining thereby the semi-direct product $C_2 \ltimes \tilde{W}^+$, where C_2 is the cyclic group of order 2. Let s be the generator of C_2. One verifies that the following maps

$$\tilde{s}_0 \mapsto s, \ \tilde{s}_i \mapsto s\tilde{R}_i, \ i = 1, \ldots, n-1, \ \alpha \mapsto z,$$

$$s \mapsto \tilde{s}_0, \ \tilde{R}_i \mapsto \tilde{s}_0\tilde{s}_i, \ i = 1, \ldots, n-1, \ z \mapsto \alpha,$$

extend to homomorphisms $\psi_1 \ : \ \tilde{G} \to C_2 \ltimes \tilde{W}^+$ and $\psi_2 \ : \ C_2 \ltimes \tilde{W}^+ \to \tilde{G}$ such that $\psi_1\psi_2 = \mathrm{Id}_{C_2 \ltimes \tilde{W}^+}$ and $\psi_2\psi_1 = \mathrm{Id}_{\tilde{G}}$. The restriction of ψ_1 to $\tilde{G}^+$ induces the homomorphism inverse to ϕ. □

Recall our convention for the orientation of edges of a Coxeter graph or of a connected extension of it: if there is an edge between vertices i and j, $i < j$, then it is oriented from i to j. Associate a generator $\tilde{r}_{ij}$ of $\tilde{G}^+$ to each generator r_{ij} of G^+. Denote by $\tilde{r}_{ji}$ the inverse of $\tilde{r}_{ij}$, $\tilde{r}_{ji} := \tilde{r}_{ij}^{-1}$. Extend the Definition 2.3 to the generators $\tilde{r}_{ij}$.

Proposition 3.2 *Let* (G, S) *be a Coxeter system with the Coxeter matrix* $\mathfrak{m}$*. The group* $\tilde{G}^+$ *is generated by* $\tilde{r}_{ij}$ *and* z *with the defining relations*

$$\begin{cases} (\tilde{r}_{ij})^{m_{ij}} = z^{m_{ij}-1} & \text{for all generators } \tilde{r}_{ij},\\ \tilde{r}_{ii_1}\tilde{r}_{i_1i_2}\ldots\tilde{r}_{i_ai} = 1 & \text{for cycles with edges } (ii_1), (i_1i_2)\ldots, (i_ai),\\ (\tilde{r}_{ij}\tilde{r}_{jk})^2 = z & \text{for } \tilde{r}_{ij}, \tilde{r}_{jk} \text{ such that } i < k \text{ and } m_{ik} = 2,\\ (\tilde{r}_{ij}\tilde{r}_{jk}\tilde{r}_{kl})^2 = z & \text{for } \tilde{r}_{ij}, \tilde{r}_{jk}, \tilde{r}_{kl} \text{ such that } i < l \text{ and } m_{il} = 2,\\ \tilde{r}_{ij}\tilde{r}_{lm} = \tilde{r}_{lm}\tilde{r}_{ij} & \text{for } \tilde{r}_{ij} \text{ and } \tilde{r}_{lm} \text{ which are not connected.}\\ z \text{ is central and } z^2 = 1. \end{cases}$$

We omit the proof as it goes along the same lines as the proof of the Proposition 2.2.

We give without details the presentations for $\tilde{G}'^{+}$. The generators of the presentation for $\tilde{G}'^{+}$ analogous to the one given in the Proposition 3.1 will be denoted by $\tilde{R}'_1, \dots, \tilde{R}'_{n-1}$ and z'. The defining relations are

$$\begin{cases} (\tilde{R}'_i)^{m_{0i}} = z' & \text{for } i = 1, \dots, n-1, \\ (\tilde{R}'^{-1}_i \tilde{R}'_j)^{m_{ij}} = z' & \text{for } i, j = 1, \dots, n-1 \text{ such that } i < j, \\ z' \text{ is central and } z'^2 = 1. & \end{cases}$$

For $\tilde{G}'^{+}$, the generators of the presentation analogous to the one given in the Proposition 3.2 will be denoted by $\tilde{r}'_{ij}$ and z' with the same conventions as above. The defining relations are

$$\begin{cases} (\tilde{r}'_{ij})^{m_{ij}} = z' & \text{for all generators } \tilde{r}'_{ij}, \\ \tilde{r}'_{ii_1} \tilde{r}'_{i_1 i_2} \dots \tilde{r}'_{i_a i} = z'^{a+1} & \text{for cycles with edges } (ii_1), (i_1 i_2) \dots, (i_a i), \\ (\tilde{r}'_{ij} \tilde{r}'_{jk})^2 = z' & \text{for } \tilde{r}'_{ij}, \tilde{r}'_{jk} \text{ such that } i < k \text{ and } m_{ik} = 2, \\ (\tilde{r}'_{ij} \tilde{r}'_{jk} \tilde{r}'_{kl})^2 = z' & \text{for } \tilde{r}'_{ij}, \tilde{r}'_{jk}, \tilde{r}'_{kl} \text{ such that } i < l \text{ and } m_{il} = 2, \\ \tilde{r}'_{ij} \tilde{r}'_{lm} = \tilde{r}'_{lm} \tilde{r}'_{ij} & \text{for } \tilde{r}'_{ij} \text{ and } \tilde{r}'_{lm} \text{ which are not connected.} \\ z' \text{ is central and } z'^2 = 1. & \end{cases}$$

Proposition 3.3 *The extensions $\tilde{G}^{+}$ and $\tilde{G}'^{+}$ of the group G^{+} are isomorphic.*

Proof. The homomorphism $\tilde{G}^{+} \to \tilde{G}'^{+}$ defined on generators by $\tilde{r}_{ij} \mapsto z' \tilde{r}'_{ij}$ and $z \mapsto z'$ establishes the required isomorphism. □

2. Fix, for type A, the numbering of the vertices and orientation of the edges as shown in Fig. 2 and set $\tilde{r}_i := \tilde{r}_{i-1,i}$, $i = 1, \dots, n-1$. The group $\tilde{A}^{+}_n$ is the spinor cover of the alternating group A^{+}_n. The presentation for $\tilde{A}^{+}_n$ from the Proposition 3.2 reads

$$\begin{cases} \tilde{r}_i^3 = 1, & \\ (\tilde{r}_i \tilde{r}_{i+1})^2 = z, & \\ (\tilde{r}_i \tilde{r}_{i+1} \tilde{r}_{i+2})^2 = z, & \\ \tilde{r}_i \tilde{r}_j = \tilde{r}_j \tilde{r}_i & \text{if } \mid i - j \mid > 2, \\ z \text{ is central and } z^2 = 1. & \end{cases} \tag{21}$$

This presentation equips the chain of the spinor extensions of the alternating groups of type A with a structure of a local and stationary tower.

3. The Schur multipliers for the alternating groups were calculated by Schur in [10]; Maxwell [6] generalized this result to alternating subgroups of finite Coxeter groups and gave a presentation, in the spirit of (2), of the corresponding central extensions. It is straightforward - although lengthy - to transform this presentation (for all the central extensions of the alternating groups of all finite Coxeter groups) to the presentation which uses generators related to the oriented edges of the Coxeter graph.

As an example we give, omitting details, the presentations for type A. The Schur multiplier of A_n^+ is C_2 if $n \geqslant 3, n \neq 5, 6$, and $C_2 \times C_3$ if $n = 5, 6$; here C_m is the cyclic group with m elements. We describe the central extensions of the groups A_5^+ and A_6^+ with the kernel $C_2 \times C_3$; these extensions are universal central extensions since each of the groups A_5^+ and A_6^+ is perfect, that is, it coincides with its commutator subgroup. As above, we associate a generator $\tilde{r}_i$, $i = 1, \dots, n-1$, to each oriented edge of the Coxeter graph, see Fig. 2.

- The universal central extension of A_5^+ is generated by $\tilde{r}_1$, $\tilde{r}_2$, $\tilde{r}_3$, $\tilde{r}_4$, z and ζ with the defining relations:

$$\begin{cases} \tilde{r}_i^3 = 1 \quad \text{for } i = 1, \dots, 4, \\ (\tilde{r}_1\tilde{r}_2)^2 = z, \quad (\tilde{r}_2\tilde{r}_3)^2 = z\zeta, \quad (\tilde{r}_3\tilde{r}_4)^2 = z, \\ (\tilde{r}_1\tilde{r}_2\tilde{r}_3)^2 = z, \quad (\tilde{r}_2\tilde{r}_3\tilde{r}_4)^2 = z, \\ \tilde{r}_1\tilde{r}_4 = \zeta^2\tilde{r}_4\tilde{r}_1, \\ z^2 = 1 \text{ and } z \text{ is central}, \quad \zeta^3 = 1 \text{ and } \zeta \text{ is central.} \end{cases} \tag{22}$$

- The universal central extension of A_6^+ is generated by $\tilde{r}_1$, $\tilde{r}_2$, $\tilde{r}_3$, $\tilde{r}_4$, $\tilde{r}_5$, z and ζ with the defining relations:

$$\begin{cases} \tilde{r}_i^3 = 1 \quad \text{for } i = 1, \dots, 5, \\ (\tilde{r}_i\tilde{r}_{i+1})^2 = z \quad \text{for } i = 1, \dots, 4, \\ (\tilde{r}_1\tilde{r}_2\tilde{r}_3)^2 = z, \quad (\tilde{r}_2\tilde{r}_3\tilde{r}_4)^2 = z\zeta, \quad (\tilde{r}_3\tilde{r}_4\tilde{r}_5)^2 = z, \\ \tilde{r}_1\tilde{r}_4 = \zeta\tilde{r}_4\tilde{r}_1, \quad \tilde{r}_2\tilde{r}_5 = \zeta\tilde{r}_5\tilde{r}_2, \quad \tilde{r}_1\tilde{r}_5 = \zeta^2\tilde{r}_5\tilde{r}_1, \\ z^2 = 1 \text{ and } z \text{ is central}, \quad \zeta^3 = 1 \text{ and } \zeta \text{ is central.} \end{cases} \tag{23}$$

4. Coxeter–Todd algorithms and normal forms for alternating groups of types A, B and D

Let G be a finite group with a given presentation by generators $g_1, \dots, g_m$ and the set of defining relations $\mathcal{R}$. Let I be a subset of $\{1, \dots, m\}$ and H the subgroup of G generated by g_a, $a \in I$. The Coxeter–Todd algorithm for the pair (G, H) constructs the set of the left cosets of H in G and the action of the generators on this set [4]. The result of a Coxeter–Todd algorithm is a figure whose vertices are labeled by left cosets and the arrows stand for the action of the generators. The algorithm starts with the left coset H; only the generators g_a, $a \notin I$, may act non-trivially on this coset and give new vertices. At each step we analyze, using the relations from $\mathcal{R}$, the action of the generators on vertices and draw new vertices or identify existing vertices. The algorithm is finished when we know the action of all generators on every coset in the figure.

The Coxeter–Todd algorithm for (G, H) lists the left cosets and thus provides a normal form for elements of G with respect to H. The algorithm implies an upper bound for the cardinality of G. Namely, let $\mathcal{H}$ be the abstract group with the generators g_a, $a \in I$; the set of defining relations for $\mathcal{H}$ is the subset of $\mathcal{R}$ consisting of those relations which involve only the generators g_a, $a \in I$. There is a natural surjection $\mathcal{H} \to H$ therefore the cardinality of G is less or equal than the numbers of vertices in the figure times the cardinality of $\mathcal{H}$. For the chain $\{1\} = G_0 \subset G_1 \subset \dots \subset G_n \subset \dots$ of groups, we construct recursively the global normal form for elements of any G_n using the normal form for elements of G_k with respect to G_{k-1}, $k = 1, 2, \dots, n$.

Here we exhibit results of the Coxeter–Todd algorithm for the chains of the alternating groups of types A, B and D with the presentation of the Proposition 2.2. As a matter of comparison we also give in each case Coxeter–Todd figures for the alternating groups with the presentation (2), for the alternating groups with a presentation à la Carmichael and for the Coxeter groups with the standard presentation (1). Each time the normal form follows straightforwardly from the Coxeter–Todd figure; we illustrate it on the example of the chain of the alternating groups of type A.

The subgroup chosen for realizing the Coxeter–Todd algorithm is always denoted by H. In a Coxeter–Todd figure, when the action of a generator g on a coset Y is not specified it means that $gY = Y$. For a generator of order 2 an unoriented edge represents a pair of oppositely oriented edges. The action of a generator of order 3 (respectively, 4) is often represented by an oriented triangle (respectively, quadrilateral); the generator is written inside it. We indicate the coset (in the form uH with u a word in the generators) at each vertex of the Coxeter–Todd figure only for type A; in general, it can be easily found following the edges from the vertex H in the figures.

4.1 Type A

The vertices and oriented edges of the Coxeter graph of type A are labeled as on Fig. 2, Section 2. The Coxeter group A_n is generated by $s_0, \dots, s_{n-1}$ with the defining relations

$$\begin{cases} s_i^2 = 1 & \text{for } i = 0, \dots, n-1, \\ s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1} & \text{for } i = 0, \dots, n-2, \\ s_i s_j = s_j s_i & \text{for } i, j = 0, \dots, n-1 \text{ such that } |i-j| > 1. \end{cases} \tag{24}$$

The group A_n is isomorphic to the group of permutations of the set $\{1, 2, \dots, n+1\}$. The isomorphism is given by $s_i \mapsto (i+1, i+2)$, $i = 0, \dots, n-1$.

Let H be the subgroup generated by $s_0, \dots, s_{n-2}$; here is the Coxeter–Todd figure for (A_n, H):



Fig. 3. *Coxeter–Todd figure for* (A_n, H) *for the presentation (24)*

We give Coxeter–Todd figures for three presentations of the alternating group A_n^+. In each situation, H is the subgroup generated by the first $n-2$ generators. The first two presentations can be found in [3]. The second one is the presentation (2). The third one is the presentation (14). To illustrate the usefulness of the Coxeter–Todd algorithm we reestablish that these are indeed presentations of the group A_n^+ and find three normal forms for elements of A_n^+.

1. In Carmichael presentation [3] the generators are $a_1, \dots, a_{n-1}$ with the defining relations

$$\begin{cases} a_i^3 = 1 & \text{for } i = 1, \dots, n-1, \\ (a_i a_j)^2 = 1 & \text{for } i, j = 1, \dots, n-1 \text{ such that } i < j. \end{cases} \tag{25}$$



Fig. 4. *Coxeter–Todd figure for* (A_n^+, H) *for Carmichael presentation (25)*

We geometrically interpret the Carmichael presentation as follows: fix the oriented edge $(0, 1)$ in the Coxeter graph of type A and take $a_1 := s_0s_1$. The other generators are obtained by consecutive conjugations of a_1 by the Coxeter generators, namely $a_i := s_ia_{i-1}s_i$, $i = 2, \ldots, n-1$. The set $\{a_1, \ldots, a_{n-1}\}$ is a generating set for the alternating group A_n^+ with the defining relations (25).

2. Moore presentation [3] is the presentation (2). The generators are $R_1, \ldots, R_{n-1}$ with the defining relations

$$\begin{cases} R_1^3 = 1, & \\ R_i^2 = 1 & \text{for } i = 2 \ldots, n-1, \\ (R_i^{-1}R_{i+1})^3 = 1 & \text{for } i = 1, \ldots, n-2, \\ (R_i^{-1}R_j)^2 = 1 & \text{for } i, j = 1, \ldots, n-1 \text{ such that } |i-j| > 1. \end{cases} \tag{26}$$



Fig. 5. *Coxeter–Todd figure for* (A_n^+, H) *for Moore presentation (26)*

3. In the presentation (14) of A_n^+ the generators are $r_1, \ldots, r_{n-1}$ with the defining relations

$$\begin{cases} r_i^3 = 1 & \text{for } i = 1, \ldots, n-1, \\ (r_ir_{i+1})^2 = 1 & \text{for } i = 1, \ldots, n-2, \\ (r_ir_{i+1}r_{i+2})^2 = 1 & \text{for } i = 1, \ldots, n-3, \\ r_ir_j = r_jr_i & \text{for } i, j = 1, \ldots, n-1 \text{ such that } |i-j| > 2. \end{cases} \tag{27}$$


H
$r_{n-1}H$
$r_{n-3}r_{n-1}H$
$r_3\dots r_{n-3}r_{n-1}H$
$r_1r_3\dots r_{n-3}r_{n-1}H$
r_{n-1}
r_{n-3}
r_1
r_{n-2}
r_2
$r^2_{n-1}H$
$r_{n-2}r^2_{n-1}H$
$r_4\dots r_{n-2}r^2_{n-1}H$
$r_2r_4\dots r_{n-2}r^2_{n-1}H$

Fig. 6. *Coxeter–Todd figure for (A_n^+, H) for the presentation (27), n even*


H
$r_{n-1}H$
$r_{n-3}r_{n-1}H$
$r_4\dots r_{n-3}r_{n-1}H$
$r_2r_4\dots r_{n-3}r_{n-1}H$
r_{n-1}
r_{n-3}
r_2
r_{n-2}
r_1
$r^2_{n-1}H$
$r_{n-2}r^2_{n-1}H$
$r_3\dots r_{n-2}r^2_{n-1}H$
$r_1r_3\dots r_{n-2}r^2_{n-1}H$

Fig. 7. *Coxeter–Todd figure for (A_n^+, H) for the presentation (27), n odd*

4. By induction on n, an upper bound, implied by the Coxeter–Todd algorithm, for the cardinality of the group defined by (25) or (26) or (27) is $n!/2$ in each case. The converse inequality is implied by the following surjective morphisms onto the group of even permutations of $n+1$ elements:

- For Carmichael presentation (25): $a_i \mapsto (1, 2, i+2)$, $i = 1, \dots, n-1$.
- For Moore presentation (26): $R_i \mapsto (1, 2)(i+1, i+2)$, $i = 1, \dots, n-1$.
- For the presentation (27): $r_i \mapsto (i, i+1, i+2)$, $i = 1, \dots, n-1$.

This concludes the proof that each of the groups - defined by (25) or (26) or (27) - has the cardinality $n!/2$ and is isomorphic to the group A_n^+ (thus the morphisms above are the isomorphisms).

For each of the three presentations, the Coxeter–Todd figure gives a list E_n of elements of A_n^+ such that any $x \in A_n^+$ can be written as $u_n h$ where $h \in A_{n-1}^+$ and $u_n \in E_n$; the set E_n is in bijection with the set of vertices of the Coxeter–Todd figure. The element h in turn can be written in a normal form with respect to A_{n-2}^+. Continuing we obtain recursively three normal forms for elements of A_n^+.

Proposition 4.1 *Let A_n^+ be given by Carmichael presentation (25). Any element $x \in A_n^+$ can be uniquely written as $x = u_n u_{n-1} \dots u_2$ where $u_i \in E_i$,*

$$E_i = \left\{1, a_{i-1}, a_{i-1}^2, a_{i-2}a_{i-1}^2, \dots, a_2 a_{i-1}^2, a_1 a_{i-1}^2\right\} \ , \ i = 2, \dots, n.$$

Proposition 4.2 *Let A_n^+ be given by Moore presentation (26). Any element $x \in A_n^+$ can be uniquely written as $x = u_n u_{n-1} \dots u_2$ where $u_i \in E_i$,*

$$E_i = \left\{1, R_{i-1}, R_{i-2}R_{i-1}, \dots, R_1R_2 \dots R_{i-1}, R_1^2R_2 \dots R_{i-1}\right\} \ , \ i = 2, \dots, n.$$

Proposition 4.3 *Let A_n^+ be given by the presentation (27). Any element $x \in A_n^+$ can be uniquely written as $x = u_n u_{n-1} \dots u_2$ where $u_i \in E_i$, $i = 2, \dots, n$, and*

- *if i is even, $E_i := \left\{1, r_{i-1}, r_{i-3}r_{i-1}, \dots, r_1r_3 \dots r_{i-3}r_{i-1}, r_{i-1}^2, r_{i-2}r_{i-1}^2, \dots, r_2r_4 \dots r_{i-2}r_{i-1}^2\right\}$,*
- *if i is odd, $E_i := \left\{1, r_{i-1}, r_{i-3}r_{i-1}, \dots, r_2r_4 \dots r_{i-3}r_{i-1}, r_{i-1}^2, r_{i-2}r_{i-1}^2, \dots, r_1r_3 \dots r_{i-2}r_{i-1}^2\right\}$.*

4.2 Type B

We label the vertices and oriented edges of the Coxeter graph of type B as follows.



Fig. 8. *Coxeter graph of type B*

The Coxeter group B_n is generated by s_0, …, s_{n-1} with the defining relations

$$\begin{cases} s_i^2 = 1 & \text{for } i = 0, \dots, n-1, \\ s_0s_1s_0s_1 = s_1s_0s_1s_0, & \\ s_is_{i+1}s_i = s_{i+1}s_is_{i+1} & \text{for } i = 1, \dots, n-2, \\ s_is_j = s_js_i & \text{for } i, j = 0, \dots, n-1 \text{ such that } |i-j| > 1. \end{cases} \tag{28}$$

The group B_n is isomorphic to the wreath product $C_2 \wr A_{n-1}$ of the cyclic group C_2 of order 2 by the symmetric group A_{n-1}. Denote by γ the generator of C_2 and by 1_{C_2} the unit element of C_2. Let $\gamma^{(i)} := (1_{C_2}, \dots, 1_{C_2}, \gamma, 1_{C_2}, \dots, 1_{C_2})$, $i = 1, \dots, n$, be the element of C_2^n with γ in position i and 1_{C_2} anywhere else; let $1_{C_2^n} := (1_{C_2}, \dots, 1_{C_2})$ denote the unit element of C_2^n and $1_{A_{n-1}}$ the unit element of A_{n-1}. The isomorphism between B_n and $C_2 \wr A_{n-1}$ is given by $s_0 \mapsto (\gamma^{(1)}, 1_{A_{n-1}})$ and $s_i \mapsto \left(1_{C_2^n}, (i, i+1)\right)$, $i = 1, \dots, n-1$.

Let H be the subgroup generated by s_0, …, s_{n-2}; here is the Coxeter–Todd figure for (B_n, H):



Fig. 9. *Coxeter–Todd figure for (B_n, H) for the presentation (28)*

This is an analogue - for the group B_n^+ - of the Carmichael presentation (the distinguished oriented edge is $(0, 1)$):

- the alternating group B_n^+ is generated by $a_1, \dots, a_{n-1}$ with the defining relations

$$\begin{cases} a_i^4 = 1 & \text{for } i = 1, \dots, n-1, \\ (a_1 a_i)^3 = 1 & \text{for } i = 2, \dots, n-1, \\ (a_1^2 a_i)^2 = 1 & \text{for } i = 2, \dots, n-1. \\ (a_1 a_i a_1 a_j)^2 = 1 & \text{for } i, j = 2, \dots, n-1 \text{ such that } i < j. \end{cases} \tag{29}$$

The presentation (2), respectively the presentation of the Proposition 2.2, reads:

- the alternating group B_n^+ is generated by $R_1, \dots, R_{n-1}$ with the defining relations

$$\begin{cases} R_1^4 = 1, & \\ R_i^2 = 1 & \text{for } i = 2, \dots, n-1, \\ (R_i^{-1} R_{i+1})^3 = 1 & \text{for } i = 1, \dots, n-2, \\ (R_i^{-1} R_j)^2 = 1 & \text{for } i, j = 1, \dots, n-1 \text{ such that } |i-j| > 1; \end{cases} \tag{30}$$

- respectively, the alternating group B_n^+ is generated by $r_1, \dots, r_{n-1}$ with the defining relations

$$\begin{cases} r_1^4 = 1, & \\ r_i^3 = 1 & \text{for } i = 2, \dots, n-1, \\ (r_i r_{i+1})^2 = 1 & \text{for } i = 1, \dots, n-2, \\ (r_i r_{i+1} r_{i+2})^2 = 1 & \text{for } i = 1, \dots, n-3, \\ r_i r_j = r_j r_i & \text{for } i, j = 1, \dots, n-1 \text{ such that } |i-j| > 2. \end{cases} \tag{31}$$

Denote by ϵ_0 the sign character of A_{n-1}. Let $\epsilon_{C_2^n} : C_2^n \to \{-1, 1\}$ be the homomorphism defined by $\epsilon_{C_2^n}(\gamma^{(i)}) = -1$ for $i = 1, \dots, n$. The alternating group B_n^+ of type B is isomorphic to the subgroup of $C_2 \wr A_{n-1}$ formed by elements (g, π), $\pi \in A_{n-1}$ and $g \in C_2^n$, such that $\epsilon_{C_2^n}(g)\epsilon_0(\pi) = 1$. The isomorphisms are given by:

- for the presentation (29), $a_i \mapsto \big(\gamma^{(1)}, (1, i+1)\big)$, $i = 1, \dots, n-1$.
- for the presentation (30), $R_i \mapsto \big(\gamma^{(1)}, (i, i+1)\big)$, $i = 1, \dots, n-1$.
- for the presentation (31), $r_1 \mapsto \big(\gamma^{(1)}, (1, 2)\big)$ and $r_i \mapsto \big(1_{C_2^n}, (i-1, i, i+1)\big)$, $i = 2, \dots, n-1$.

Let H be the subgroup generated by $a_1, \dots, a_{n-2}$, or by $R_1, \dots, R_{n-2}$, or by $r_1, \dots, r_{n-2}$.



Fig. 10. *Coxeter–Todd figure for* (B_n^+, H) *for the presentation (29)*



Fig. 11. *Coxeter–Todd figure for* (B_n^+, H) *for the presentation (30)*



Fig. 12. *Coxeter–Todd figure for* (B_n^+, H) *for the presentation (31), n even*



Fig. 13. *Coxeter–Todd figure for* (B_n^+, H) *for the presentation (31), n odd*

As for type A, the Coxeter–Todd figures for the presentations (29), (30) and (31) provide three normal forms for elements of the alternating group B_n^+. We omit details.

Remark. The complex reflection group $G(m,1,n)$ is generated by s_0, …, s_{n-1} with the same defining relations as in (28), except that s_0^2 is replaced by $s_0^m = 1$. For $m = 1$, $G(1,1,n)$ is the group A_{n-1}; for $m = 2$, $G(2,1,n)$ is the group B_n. Let H be the subgroup of $G(m,1,n)$ generated by s_0, …, s_{n-2}. The Coxeter-Todd figure for $\big(G(m,1,n), H\big)$, see [9], generalizes Fig. 3 and Fig. 9.

4.3 Type D

We label the vertices and oriented edges of the Coxeter graph of type D as follows.



Fig. 14. *Coxeter graph of type D*

The Coxeter group D_n is generated by $s_0, \dots, s_{n-1}$ with the defining relations

$$\begin{cases} s_i^2 = 1 \quad \text{for } i = 0, \dots, n-1, \\ s_0 s_2 s_0 = s_2 s_0 s_2, \quad s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1} \quad \text{for } i = 1, \dots, n-2, \\ s_0 s_1 = s_1 s_0, \quad s_0 s_i = s_i s_0 \quad \text{for } i = 3, \dots, n-1, \\ s_i s_j = s_j s_i \quad \text{for } i, j = 1, \dots, n-1 \text{ such that } |i-j| > 1. \end{cases} \tag{32}$$

The group D_n is isomorphic to the subgroup of $C_2 \wr A_{n-1}$ formed by elements (g, π), $\pi \in A_{n-1}$ and $g \in C_2^n$, such that $\epsilon_{C_2^n}(g) = 1$. The isomorphism is given by $s_0 \mapsto \big(\gamma^{(1)}\gamma^{(2)}, (1,2)\big)$ and $s_i \mapsto \big(1_{C_2^n}, (i, i+1)\big)$ for $i = 1, \dots, n-1$.

Let H be the subgroup generated by $s_0, \dots, s_{n-2}$; here is the Coxeter–Todd figure for (D_n, H):



Fig. 15. *Coxeter–Todd figure for (D_n, H) for the presentation (32)*

This is an analogue - for the group D_n^+ with $n \geq 3$ - of the Carmichael presentation (the distinguished oriented edge is $(0, 2)$):

- the alternating group D_n^+, $n \geq 3$, is generated by $a_1, \dots, a_{n-1}$ with the defining relations

$$\begin{cases} a_i^3 = 1 & \text{for } i = 1, \dots, n-1, \\ (a_1 a_i)^2 = 1 & \text{for } i = 2, \dots, n-1, \\ (a_2^2 a_i)^2 = 1 & \text{for } i = 3, \dots, n-1, \\ (a_i a_j)^2 = 1 & \text{for } i, j = 3, \dots, n-1 \text{ such that } i < j. \end{cases} \tag{33}$$

The presentation (2), respectively the presentation of the Proposition 2.2, reads:

- the alternating group D_n^+ is generated by $R_1, \ldots, R_{n-1}$ with the defining relations

$$\begin{cases} R_2^3 = 1, & \\ R_i^2 = 1 & \text{for } i = 1, \ldots, n-1 \text{ and } i \neq 2, \\ (R_i^{-1}R_{i+1})^3 = 1 & \text{for } i = 1, \ldots, n-2, \\ (R_i^{-1}R_j)^2 = 1 & \text{for } i, j = 1, \ldots, n-1 \text{ such that } |i-j| > 1; \end{cases} \tag{34}$$

- respectively, the alternating group D_n^+, $n \geq 3$, is generated by $r_1, \ldots, r_{n-1}$ with the defining relations

$$\begin{cases} r_i^3 = 1 \quad \text{for } i = 1, \ldots, n-1, \\ (r_1r_2^2)^2 = 1, \quad (r_1r_3)^2 = 1, \quad (r_ir_{i+1})^2 = 1 \quad \text{for } i = 2, \ldots, n-2, \\ (r_1r_3r_4)^2 = 1, \quad (r_ir_{i+1}r_{i+2})^2 = 1 \quad \text{for } i = 2, \ldots, n-3, \\ r_1r_i = r_ir_1 \quad \text{for } i = 5, \ldots, n-1, \\ r_ir_j = r_jr_i \quad \text{for } i, j = 2, \ldots, n-1 \text{ such that } |i-j| > 2. \end{cases} \tag{35}$$

The alternating group D_n^+ of type D is isomorphic to the subgroup of $C_2 \wr A_{n-1}$ formed by elements (g, π), $\pi \in A_{n-1}$ and $g \in C_2^n$, such that $\epsilon_{C_2^n}(g) = 1$ and $\epsilon_0(\pi) = 1$. The isomorphisms are given by:

- for the presentation (33),
  $a_1 \mapsto \big(\gamma^{(1)}\gamma^{(2)}, (1,2,3)\big)$, $a_2 \mapsto \big(\gamma^{(1)}\gamma^{(2)}, (1,3,2)\big)$ and $a_i \mapsto \big(\gamma^{(1)}\gamma^{(2)}, (1,2,i+1)\big)$, $i = 3, \ldots, n-1$.
- for the presentation (34),
  $R_1 \mapsto \big(\gamma^{(1)}\gamma^{(2)}, 1_{A_{n-1}}\big)$ and $R_i \mapsto \big(\gamma^{(1)}\gamma^{(2)}, (1,2)(i,i+1)\big)$, $i = 2, \ldots, n-1$.
- for the presentation (35),
  $r_1 \mapsto \big(1_{C_2^n}, (1,2,3)\big)$, $r_2 \mapsto \big(\gamma^{(1)}\gamma^{(2)}, (1,2,3)\big)$ and $r_i \mapsto \big(1_{C_2^n}, (i-1,i,i+1)\big)$, $i = 3, \ldots, n-1$.

Let H be the subgroup generated by $a_1, \ldots, a_{n-2}$, or by $R_1, \ldots, R_{n-2}$, or by $r_1, \ldots, r_{n-2}$.



Fig. 16. *Coxeter–Todd figure for (D_n^+, H) for the presentation (33)*



Fig. 17. *Coxeter–Todd figure for* (D_n^+, H) *for the presentation (34)*



Fig. 18. *Coxeter–Todd figure for* (D_n^+, H) *for the presentation (35), n even*



Fig. 19. *Coxeter–Todd figure for* (D_n^+, H) *for the presentation (35), n odd*

The Coxeter–Todd figures for the presentations (33), (34) and (35) provide three normal forms for elements of the alternating group D_n^+. We omit details.

Acknowledgements. We thank I. Marin, J. Michel, O. Brunat, C. Blanchet for useful discussions.

References

[1] Bourbaki N., *Groupes et algebres de Lie*, Springer (2002).

[2] Brenti F., Reiner V. and Roichman Y., *Alternating subgroups of Coxeter groups*, J. Combin. Theo. A 115(5) (2008) 845–877.

[3] Coxeter H. and Moser W., *Generators and relations for discrete groups*, Springer-Verlag, 1980.

[4] Coxeter H. and Todd J., *A practical method for enumerating cosets of a finite abstract group*, Proc. Edimburgh Math. Soc. 5 (1936) 26–34.

[5] Ihara S. and Yokonuma T., *On the second cohomology groups (Schur multipliers) of finite reflection groups*, J. Fac. Sci. Univ. Tokyo Sect. I (1965) 155–171.

[6] Maxwell G., *The Schur multipliers of rotation subgroups of Coxeter groups*, J. Algebra 53 (1978) 440–451.

[7] Morris A., *Projective representations of reflection groups*, Proc. London Mat. Soc. 32(3) (1976) 403–410.

[8] Okounkov A. and Vershik A., *A new approach to representation theory of symmetric groups II*, Selecta Math. New series 2(4) (1996) 581–605.

[9] Ogievetsky O. and Poulain d'Andecy L., *Cyclotomic Hecke algebras: Jucys-Murphy elements, representations, classical limit*, hal-0064782.

[10] Schur I., *Über die Darstellung der symmetrischen und der alternierenden Gruppe durch gebrochene lineare Substitutionen*, J. Reine Angew. Math. 139 (1911) 155–250.

[11] Vershik A., *Local stationary algebras*, Amer. Math. Soc. Trans. 148(2) (1991) 1–13.

[12] Vershik A. and Vsemirnov M., *The local stationary presentation of the alternating groups and the normal form*, J. Algebra 319(10) (2008) 4222–4229.

[13] Yokonuma T., *On the second cohomology groups (Schur multipliers) of infinite discrete reflection groups*, J. Fac. Sci. Univ. Tokyo Sect. I (1965) 173–186.